



SEKOLAH TINGGI PERTANIAN KUTAI TIMUR

Jl. Soekarno-Hatta No. 01 Sangatta, Kab. Kutai Timur Kalimantan Timur, Kode Pos 75387
Telp. (0549) 2031985 E-mail : stiper_kutim@yahoo.com

LEMBAR HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW KARYA ILMIAH : JURNAL ILMIAH BEREPUTASI TERINDEK SCOPUS

Judul Jurnal Ilmiah (Artikel) : Structure of Reef Fish Assemblages at Miang Island, East Kalimantan (Borneo)
Penulis Jurnal Ilmiah : Rosdianto (Penulis 1) dari 5 Penulis
Identitas Jurnal Ilmiah : a. Nama Jurnal : Journal of Hunan University (Natural Sciences)
b. Volume/Nomor : 48 No. 9 (2021)
c. Edisi (Bulan/Tahun) : September, 30,2021/Issue/view/78
d. Penerbit : The Journal of Hunan University Natural Sciences publishes special issues on various and relevant topics of interest to the scientific community ISSN 1674-2974
e. Jumlah Halaman : 233 - 240

Kategori Publikasi Karya Ilmiah :
(beri ✓ pada kategori yang tepat)

Jurnal Ilmiah Internasional / Internasional Bereputasi
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional Tidak Terakreditasi

Hasil Penilaian Peer Review : Artikel ditulis dengan mengikuti kaidah jurnal ilmiah internasional bereputasi. Sistematisnya jelas dan terpilah-pilah dengan baik. Research gap dipaparkan dengan baik dan menjadi alasan yang kuat untuk memunculkan penelitian. Ruang lingkup pembahasannya proporsional dan dikaji secara mendalam, sesuai dgn bidang ilmu Penulis. Hal yang patut dibanggakan adalah tulisan ini diterbitkan oleh The Journal of Hunan University Natural Sciences publishes special issues on various and relevant topics of interest to the scientific community, yang terindeks pada jurnal Internasional bereputasi, yakni Scopus dan Scimagojr (Q2).

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah			Nilai Akhir Yang Diperoleh
	Internasional/ Int. Bereputasi	Nasional Terakreditasi	Nasional Tidak Terakreditasi	
a. Kelengkapan Unsur Isi Artikel (10%)	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	2,39
b. Ruang Lingkup & Kedalaman Pembahasan (30%)	4	<input type="checkbox"/>	<input type="checkbox"/>	7,1
c. Kecukupan & Kemutakhiran Data/Informasi & Metodologi (30%)	12	<input type="checkbox"/>	<input type="checkbox"/>	7,1
d. Kelengkapan Unsur & Kualitas Terbitan/Jurnal (30%)	12	<input type="checkbox"/>	<input type="checkbox"/>	7,2
Total = (100%)	40	<input type="checkbox"/>	<input type="checkbox"/>	23,79

Sangatta, 5 Oktober 2021
Reviewer 1

Prof. Dr. Ir. Mulyono S. Baskoro, MSc
NIP : 19620303198803 1 001
Jabatan : Guru Besar 1050 / IVE
Unit Kerja : Dep. PSP, FPIK, IPB Bogor.



SEKOLAH TINGGI PERTANIAN KUTAI TIMUR

Jl. Soekarno-Hatta No. 01 Sangatta, Kab. Kutai Timur Kalimantan Timur, Kode Pos 75387
Telp. (0549) 2031985 E-mail : stiper_kutim@yahoo.com

LEMBAR HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW KARYA ILMIAH : JURNAL ILMIAH BEREPUTASI TERINDEK SCOPUS

- Judul Jurnal Ilmiah (Artikel) : **Structure of Reef Fish Assemblages at Miang Island, East Kalimantan (Borneo)**
- Penulis Jurnal Ilmiah : Rosdianto (Penulis 1) dari 5 Penulis
- Identitas Jurnal Ilmiah :
- a. Nama Jurnal : Journal of Hunan University (Natural Sciences)
 - b. Volume/Nomor: 48 No. 9 (2021)
 - c. Edisi (Bulan/Tahun): September, 30,2021/Issue/view/78
 - d. Penerbit : The Journal of Hunan University Natural Sciences publishes special issues on various and relevant topics of interest to the scientific community **ISSN 1674-2974**
 - e. Jumlah Halaman: 233 - 240
- Kategori Publikasi Karya Ilmiah : Jurnal Ilmiah Internasional / Internasional Bereputasi
(beri ✓ pada kategori yang tepat) Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional Tidak Terakreditasi

Hasil Penilaian Peer Review : Kelengkapan Isi artikel sangat baik dan memuaskan. sistematik penulisan jelas dan baik. Artikel berasal dari riset yang dilakukan secara teliti dan detail. Sistematika penulisan terjalin dengan rapi. Ruang lingkup pembahasan Sangat kuat dalam konteks kelautanan sesuai dgn bidang ilmu Penulis. Data lengkap dan mutahir dan metodologi yg digunakan sudah tepat. Hal yang patut dibanggakan adalah tulisan ini diterbitkan oleh The Journal of Hunan University Natural Sciences publishes special issues on various and relevant topics of interest to the scientific community yang terindeks pada jurnal Internasional bereputasi, yakni Scopus dan Scimagojr (Q2)/SJR 026.

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah			Nilai Akhir Yang Diperoleh
	Internasional/ Int. Bereputasi	Nasional Terakreditasi	Nasional Tidak Terakreditasi	
	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
a. Kelengkapan Unsur Isi Artikel (10%)	4			2,4
b. Ruang Lingkup & Kedalaman Pembahasan (30%)	12			7,2
c. Kecukupan & Kemutahiran Data/Informasi & Metodologi (30%)	12			7,2
d. Kelengkapan Unsur & Kualitas Terbitan/Jurnal (30%)	12			7,1
Total = (100%)	40			23,9

Sangatta, 7 Oktober 2021

Reviewer 2



Prof. Dr. Ir. Juraemi, M.Si.
NIP. 195704131987021001

Jabatan Fungsional : Guru Besar 850
Unit Kerja : Rektor/Ketua STIPER



Home > Journal of Hunan University Natural Sciences

Journal of Hunan University Natural Sciences



The Journal of Hunan University Natural Sciences is the leading Chinese academic journal that publishes articles in all areas of natural sciences. The Journal is meant to serve as a means of communication and discussion of important issues related to science and scientific activities. The Journal publishes only original articles in English which have international importance. In addition to full-length research articles, the Journal publishes review articles. Papers can be focused on fundamental research leading to new methods, or adaptation of existing methods for new applications.

Articles for the Journal are peer-reviewed by third-party reviewers who are selected from among specialists in the subject matter of peer-reviewed materials.

The Journal of Hunan University Natural Sciences is a kind of forum for discussing issues and problems facing science and scholars, as well as an effective means of interaction between the members of the academic community. The Journal of Hunan University Natural Sciences is read by a large number of scholars, and the circulation of the journal is constantly growing.

The Journal of Hunan University Natural Sciences publishes special issues on various and relevant topics of interest to the scientific community.

The Journal of Hunan University Natural Sciences is indexed by Web of Science, Scopus, Current Contents, Geobase and Chemical Abstracts.

Articles containing fundamental or applied scientific results in all areas of the natural sciences are accepted for consideration.

The Editorial Board of the Journal of Hunan University Natural Sciences is composed of 25 members and is chaired by Academician Chen Zhengqing. Editor-in-chief is Prof. Yi Weijian.

Frequency of publication: monthly

ISSN: 1674-2974

Access to all articles on the website is open, does not require registration or payment.

Journal articles are licensed under the CC BY 4.0 [Creative Commons Attribution 4.0 License](#).

The Journal of Hunan University Natural Sciences takes care of maintaining electronic versions of articles. Data safety is ensured by backing up digital data in accordance with internal regulations. Logical and physical data migration is also provided. Cloud technologies are applied.

For further information, please contact:

E-mail: editorial-office@jonuns.com

Address: Lushan Road (S), Yuelu District, Changsha, Hunan Province, Zip Code: 410082 (Editorial Department of Journal)

Paper Submission

Vol 48, No 9 (2021)

Table of Contents

Articles

- | | |
|--|-----|
| A Review of Meaningful Learning through Virtual Reality Learning Environment
<i>Ghaliya Al Farsi, Azmi Bin Mohd. Yusof, Mohd Ezanee Bin Rusli</i> | PDF |
| Adoption of IoT-based Smart Healthcare: An Empirical Analysis In the Context of Pakistan
<i>Zulfiqar Ali Solangi, Yasir Ali Solangi, Zulfikar Ahmed Maher</i> | PDF |
| Analysis of Students' Mathematical Problem-Solving Skills
<i>Intan Kemala Sari, Ahmad Nasriadi, Riza Agustiani, Rahmat Nursalim, Jarudin</i> | PDF |
| Analysis of the Audiovisual Media Development for Basic Self-defense Learning in Sports Science Colleges
<i>Agus Mukholid</i> | PDF |
| Assessment of Surface Roughness and Compressive Strength of Resin-Based Composite Using Different Types of Bleaching Protocols
<i>Hemn Muhssin Suleman, Diyar Khalid Bakr, Bassam Karem Amin, Sazan Sherdil Salem</i> | PDF |
| Comparative Study of Some New Naproxen Derivatives on the Liver and Kidney Tissues of Mice
<i>Sadiq Al-Mansury, Nabaa Hameed Chekhyor, Adnan M. Jassim, Marwah Najeh Hammud, Saraa Amaid Kadium, Asaad H. Enad</i> | PDF |
| Complete Recovery Chronic Retro-Odontoid Pseudotumor Et HNP with Laminectomy and Cervical Disc Replacement
<i>I. Nyoman Semita, Yuna Annisa Salsabila, Ni Njoman Juliasih</i> | PDF |
| Correlation between Neutrophil-to-Lymphocyte Ratio with Disease Severity in Diabetic Patients with COVID-19 at Tertiary Referral Hospital in Indonesia
<i>Cupuwatie Cahyani, Hermina Novida, Soebagijo Adi Soelistijo, Usman Hadi, Nenci Siagian</i> | PDF |

- Digital Technologies as Marketing Tools to Promote Commerce in Micro and Small Businesses in Tulancingo de Bravo, Hidalgo, Mexico** PDF
Liliana de Jesús Gordillo Benavente, Benedicta María Domínguez Valdez, Claudia Vega Hernández, Edwin Chofo Ortega Cruz
- Does Herd Behavior Make the Market More Efficient?** PDF
Fitri Ismiyanti, Putu Anom Mahadwartha, Zunairoh, Santi Novita
- Early Prediction of Students' Academic Achievement: Categorical Data from Fully Online Learning on Machine-Learning Classification Algorithms** PDF
Tuti Purwoningsih, Harry B. Santoso, Kristanti A. Puspitasari, Zainal A. Hasibuan
- Effect of Glutamine Enteral Supplementation in Post-Operative Intestinal Obstruction Neonates: A Randomized Control Trial** PDF
Martono Tri Utomo, Mahendra Tri Arif Sampurna, Muhammad Pradhika Mapindra, Muhammad Pradhiki Mahindra, Setya Mirtha, Kartika Dharma Handayani, Dina Angelika, Risa Etika
- Effects of the Sonchus Arvensis Fraction in Rats with Monosodium Urate-Induced Gouty Arthritis** PDF
Nita Parisa, Rachmat Hidayat, Ziske Maritska, Bintang Arroyantri Prananjaya
- Evaluating the Stability of the Fractured Bone Implanted with Titanium Elastic Nails in C and S Configurations** PDF
Abdolreza Rastitalab, Salar Khajepour, Javad Dehghani, Ahmad Afsari, Shahin Heidari
- Evaluation of Rice Farmers' Climate Field School with "Context, Input, Process, and Product" Model** PDF
Lukman Effendy, Yayon Haryanto
- Experimental Textbook for Technical and Natural Science Subjects in the Slovak Republic** PDF
Lubomír Žáčok, Renáta Bernátová, Milan Bernát
- Histological and Biochemical Evaluation of the Effect of Desloratadine Drug in Hepatic Tissues** PDF
E.J. Jabber, E.R. Sarhat, S.A. Wadee, B.I. Sedeeq, T.R. Sarhat
- Immunohistochemical Expression of Galectin-9 in Oral Squamous Cell Carcinoma** PDF
Sehrish Ahmed, Fouzia Shaikh, Faraz Ahmed Baig, Rehan Ahmed Siddiqui, Akhtar Nasir Jamal Baig
- Improvement of Knowledge through Effective Communication Training and Soft Skills** PDF
Rita Kartika Sari
- Improving Marketing Performance of Small-Medium Enterprises of Food Stall by Innovation on Online Orders** PDF
*Widodo **
- Inclusion School for Early Childhood in the Outdoor Classroom** PDF
Agung Prasetyo, Sunardi, Nunuk Suryani, Asrowi
- Internal Fertilization of Oviparous Fishes, Oryzias Celebensis: Is It a Coincidence?** PDF
Khusnul Yaqin, Sri Wahyuni Rahim, Dwi Kesuma Sari
- Lapin Immune Features of Experimental Escherichia Coli-Pseudomonas Aeruginosa Combined Bacterin: A Research Account** PDF
Ibrahim M.S. Shnawa

- Life History Strategies of Crustose Coralline Algae (Rhodophyta) as Resilience Indicator of Coral Reef at Nature Reserve Pulau Sempu** PDF
Andik Isdianto, Oktiyas Muzaky Luthfi, Guntur, Muhammad Arif Asadi, Nuddin Harahab, Andi Kurniawan, Agus Dwi Wicaksono, Rosdianto, Berlania Mahardika Putri
- Machine Learning Predictions for the Advancement of the Online Education in the Higher Education Institutions in Jordan** PDF
Bayan Alfayoumi, Mohammad Alshraideh, Martin Leiner, Iyad Muhsen AlDajani
- Management Information Technology for Depression Strategies during COVID-19 Using by Data Mining Technique** PDF
Kanakarn Phanniphong, Praty Nuankaew
- Mass Media and Public Attitudes towards Compliance with the Implementation of Social Policies and Physical Distancing** PDF
Noorhidayah, Elsi Setiandari Lo
- Polyphenolics and Antioxidant Potential of Five Medicinal Plants Found in East Kalimantan, Indonesia** PDF
Anton Rahmadi, Miftakhur Rohmah, Kartika Sari, Medi Hendra, Riris Novia Azemi, Miftakhul Rahmadani Alfitriyah, Fatchur Rohman
- Potential CRISPR/Cas9 Associated Lenti-sgR5-Cas9 for CCR5 and CXCR4 Disruption Protects CD4+T** PDF
Nurul Hidayah, Totalenesya Reforment Sutikno, Yuna Annisa Salsabila
- Potential Cytotoxic, Antifungal, and Antioxidant Activity of Dithymoquinone and Thymoquinone** PDF
Eman Ramadan Elsharkawy, Emad M. Abdallah, Ahmad Abo Markb
- Psychoeducation Dhikr Influence on Oxytocin and Interleukin-8 Levels on Pregnant Women** PDF
Sri Wahyuni, Emy Suryani, Sri Sumarni, Henik Istikhomah, Murwati
- Restorative Justice and Its Connection with the Tolerance of the Islamic Religion in Saudi Arabia** PDF
Sallim Hamid N. Almjnoni, Mohamad Rizal Bin Abd Rahman
- South Halmahera District Government Policy toward Access to Anglers' Group Marketing in the District of the Jouronga Islands** PDF
Mohbir Umasugi, Muhammad Darsan Hi Adam
- Structure of Reef Fish Assemblages at Miang Island, East Kalimantan (Borneo)** PDF
Rosdianto, Oktiyas Muzaky Luthfi, Dhira Khurniawan Saputra, Wisnu Arditya Maulana, Andik Isdianto, Muhammad Arif Asadi
- The Application of Game-and-Simulation-Based Online Learning in Elementary School** PDF
A. Jaya, Y. Sunarya, M. Supriatna, D.S. Fuadi, W.R.M.I. Wahyono
- The Causal Relationship between Electric Power Consumption and Economic Growth in Malaysia and Thailand: ARDL Bound Testing Approach** PDF
Tanattrin Bunnag
- The Correlation between Social Media Addiction and Life Satisfaction among University Students** PDF
Adel Eladl, Ali Al Musawi

The Development of a Newly Formed Superior Local Chicken in Indonesia – A Review

PDF

Bachtar Bakrie, Eni Siti Rohaeni, Yenny Yusriani, Siska Tirajoh

The Development of Work-Based Learning and Group Investigation Learning Models in Vocational Education Diploma III Nursing

PDF

R. Taufiq Nur Muftiyanto, Muhammad Akhyar, Munawir Yusuf, Sri Haryati

The Effectiveness of Using GeoGebra Software in Mathematics Classrooms: A Case Study of Teaching Continuous Functions in High Schools

PDF

Duong Huu Tong, Bui Phuong Uyen, Huynh Thi Diem Kieu, Lu Kim Ngan

The Graceful Labeling on W Graph

PDF

Wed Giyarti, Kiki A. Sugeng, Fery Firmansah

The Impact of Energy Drinks on Surface Roughness, Hardness, and Color Stability of Three Types of Composite Restorations

PDF

D.I. Hamadamin, D.H. Saeed

The Influence of Balanced Scorecard on Management Accounting Performance in Malaysia during the Covid-19 Pandemic

PDF

Sudhashini Nair, Lee Yong Hunt, Neeta Jayabalan

The Needs Analysis of the Instructional Media to Improve Students' Entrepreneur Motivation in Historical Academic Teaching

PDF

Soebijantoro, Furqon Hidayatullah, Samsi Haryanto, Sri Haryati

The Role of Market Fluctuation and Imperfection on Dividends Smoothing

PDF

Masoud Taherinia, Abas Talari

Tor Attack Studies: A Survey on Active and Passive Subject

PDF

Mohamad Amar Irsyad Mohd Aminuddin, Zarul Fitri Zaaba, Azman Samsudin, Nor Badrul Anuar Juma'at



- Editorial Team
- Editorial Policies
- Focus & Scope
- Author Guidelines
- Abstracting and Indexing
- Publication Ethics
- Paper Submission

INDEXED BY SCOPUS

Open Access Article

Structure of Reef Fish Assemblages at Miang Island, East Kalimantan (Borneo)

Rosdianto^{1*}, Oktiyas Muzaky Luthfi², Dhira Khurniawan Saputra², Wisnu Arditya Maulana²,
Andik Isdianto², Muhammad Arif Asadi²

¹ Marine Science, East Kutai Agricultural College School (STIPER), Jl. Soekarno Hatta, Tlk. Lingga, Sangatta, Kabupaten Kutai Timur, Kalimantan Timur 75683, Indonesia

² Department of Marine Sciences, Faculty of Fisheries and Marine Sciences, University of Brawijaya, Jl. Veteran, Malang 65145, Indonesia

Abstract: Miang Island (118°0'20E-0°44'0N), located in East Kutai Regency, East Kalimantan, has an island area of 7.39 km² with a coastline length of about 11.73 km². Coral reef ecosystems can be found around the island by forming a fringing reef with an area of about ± 218.8 ha. The existence of this reef ecosystem certainly attracts a variety of biota, including reef fish. The abundance of reef fish in Miang Island has not been proven because no previous research has discussed reef fish in this region. Prioritizing conservation areas requires knowledge of the spatial distribution of fish assemblages' richness and organization. This research describes the biodiversity and community structure of reef fish assemblages in Pulau Miang at two different depths. The Underwater Visual Census (UVC) has been used for reef fish data retrieval. The results showed the abundance of reef fish at station 1 amounted to 0.45 ind./m³, station two by 0.10 ind./m³, station three by 0.06 ind./m³, and station four by 0.09 ind./m³. A total of 88 species from 21 families were successfully described in this study. The abundance of reef fish is dominated by the families Pomacentridae and Caesionidae, as these two families are temperate fish, which are often found in tropical waters. Our findings may be important as primer data for managers to plan this region's future marine protected area.

Keywords: Pomacentridae, Caesionidae, habitat, competition, reef fish, remote area.

東加里曼丹島米昂島礁魚群結構 (婆羅洲)

摘要: 要米昂島 (118° 0'20 乙-0° 44'0 N) 位於東庫台麗晶, 東加里曼丹島面積 7.39 公里 2, 海岸線長約 11.73 公里 2。珊瑚礁生態系統可以通過形成面積約 ±218.8 公頃的珊瑚礁來發現。這個珊瑚礁生態系統的存在當然吸引了各種生物群落, 包括珊瑚礁魚。米昂島珊瑚礁魚類的豐度尚未得到證實, 因為以前沒有詳細討論過該地區珊瑚礁魚類的存在。優先保護地區需要了解魚類組合的豐富性和組織性的空間分佈。這項研究的目的是描述普勞米昂珊瑚礁魚群的生物多樣性和社區結構在兩個不同的深度。水下視覺普查 (紫外線) 已用於珊瑚礁魚類數據檢索。結果顯示, 第 1 站礁魚的豐度為 0.45 人/立方米, 第 2 站為 0.10 人/立方米, 第 3 站為 0.06 人/立方米, 第 4 站為 0.09 人/米 3。這項研究共成功描述了來自 21 個家庭的 88 種物種。珊瑚礁魚的豐度由波馬森特裡達和凱索尼達家族主導, 因為這兩個家庭是溫帶魚, 經常在熱帶水域發現。

关键词: 果蠅科, 彩蝶科, 栖息地, 竞争, 礁鱼, 偏远地区。

1. Introduction

Received: May 1, 2021 / Revised: June 6, 2021 / Accepted: August 1, 2021 / Published: September 30, 2021

About the authors: Rosdianto, Marine Science, East Kutai Agricultural College School (STIPER), Sangatta, Indonesia; Oktiyas Muzaky Luthfi, Dhira Khurniawan Saputra, Wisnu Arditya Maulana, Andik Isdianto, Muhammad Arif Asadi, Department of Marine Sciences, Faculty of Fisheries and Marine Sciences, University of Brawijaya, Malang, Indonesia

Corresponding author Rosdianto, rosdianto73anto@gmail.com

Diversity, in general, can be determined as a condition of spreading or composing biota evenly throughout the earth, although various biota or living things are more likely to be concentrated somewhere [1]. This has happened in reef fish, which are concentrated in the coral triangle region. As many as 37% of reef fish worldwide are found in Indonesia and the Philippines. Indonesia has the highest diversity of reef fish species globally, with more than 2500 reef fish species recorded [2] and is estimated to increase to 2600 species [3]. Furthermore, there will be various hypotheses why the Coral Triangle Region, the other term is Indo-Australian Archipelago (IAA), to be the center of reef fish biodiversity?

The first hypothesis states that the IAA Region is the center of origin, which are all come from here [4]. Second, the center of overlap is a consequence of the intersection between the Indian and Pacific Oceans [5]. Third, the center of accumulation, where the number of corals that live in this aquatic area attracts various biota, including reef fish [6, 7].

Reef fish have two active roles in coral reef ecosystems. First, they can be an indicator of the health of coral reef ecosystems [8], and second, they can be used to measure the level of habitat suitability because reef fish move around to choose habitats with more suitable conditions for their life [9].

As well known that herbivorous fish can control the growth of algae (turf algae, corticated algae, tough erect algae, and foliose algae) and give a chance for coral to occupy the substrate and grow fast [10, 11]. Another function of reef fish is the certain types of reef fish in a coral reef area were accurate instructions to describe the health condition of the ecosystem. For instance, the presence of the family *Chaetodontidae*, whose life depends on the existence of corals, so the existence of this fish is usually used as a bioindicator of the health of coral reefs in general [12, 13].

Pulau Miang (118°0'20E-0°44'0N) administratively located at Sangkulirang district, East Kutai Regency, East Kalimantan. This island has about a 7.39 km² area with a coastline length of about 11.73 km [14]. Pulau Miang is the name of two islands they are Pulau Miang Kecil and Pulau Miang Besar. Pulau Miang Kecil is not an inhabited island, the wide-area was 0.82 km², and the coastline was 3.6 km. Meanwhile, Pulau Miang Besar was inhabited by 500 people that were fully dependent on fisheries activities. When local people talk about Pulau Miang, it will refer to Pulau Miang Besar, used in this research.

A fringing reef circled Pulau Miang, and in the north-western part of this island was spread some patch reef [14]. The biodiversity and potential of coral reefs on this island have not been discussed yet, since the location was in a remote area, and the high accommodation cost can be two reasons why fewer reports regarding the coral reef biodiversity on this island.

The purpose of this research is to provide an overview of the condition and biodiversity of the reef fish located on Pulau Miang as the basis of future conservation policies by the local government.

2. Materials and Methods

2.1. Study Site

This research was conducted in Pulau Miang Besar (Pulau Miang), East Kalimantan on April 28 – May 3, 2018. This location was chosen because the island has the best coral reef cover area in Sangkulirang District and is one of the best in East Kutai Regency [14]. The research location was conducted at four stations that were represented of the northern part (St.1), the western part (St.2), the eastern part (St.3), southern part (St.4) of Pulau Miang. Localities of each station were chosen by randomly based on information of local people and there was coordinate each stations: station 1 (00° 58.816' N, 117° 58.538' E); station 2 (00°43.839' N, 117°59.551' E); station 3 (00°43.901' N, 118°01.385' E); and station 4 (00°43.044' N, 118°01.172' E) (Fig. 1).




Fig. 1 The localities of the research station were marked by the blue point

2.2. Distribution of Reef Fish

This study's reef fish data collection used the UVC method (Underwater Visual Census) [15]. The 100 m-long transects have lied parallel to the shoreline in five and ten m depth than two observers were dive into right and left of the transect calculated and record the reef fish on 2.5 m right and left of transect using an underwater video camera (Canon G-16, Japan). Observer also records fish five m above transect along of transect that resulted in 100x5x5 m (2500 m³). While recording the underwater video camera, observers also calculated the number of fish on their area in the underwater slate and estimated the fish size. All reef fish data was conducted on five and ten m depth following the contours of coral in Pulau Miang than found on those depths. Observers also photographed the reef fish to obtain good quality fish pictures for identification. All data from the underwater camera was then transferred into a computer and interpreted the fish data to an Excel file.




Fig. 2 Illustration of reef fish data retrieval. Fish identification

Identification of the genus of reef fish was carried out directly in the field and backed up using underwater photo documentation. The fish that had been recognized are recorded directly on the underwater slate, while some types of fish that have not been identified will be identified referring to the manual after the data retrieval is complete—identification of the species of reef fish used protocol by [2, 16].

2.3. Data Analysis

The abundance of the number of fish found per unit of the transect was calculated using the formula that derived as:

$$X_i = n_i/A \quad (1)$$

where:

X_i = Abundance of i -type fish (ind./m³)

n_i = Number of i -type fish

A = Area of observation

Reef fish abundance and distribution data were analyzed descriptively using charts and tables in Microsoft Office Excel 2010 and expressed as a numerical individual per meter square [8, 17].

3. Results and Discussion

3.1. Reef Fish in Pulau Miang

The results of this research observed in total about 704 reef fish from 21 different families. The distribution of reef fish in shallow water (5 m depth) was less compared to deep water (10 m depth), in which the population was 296 and 408 fish for each depth (Fig. 3). As shown in Fig. 3 and Table 1, the population of Pomacentridae in Pulau Miang was the highest among others. Pomacentridae is unique reef fish that have territorial in the coral reef ecosystem, which had structured the benthic communities on their territory and mammals in the land territory [18]. Through their feeding habit and behavior activities, pomacentrids fish mediated the growth of algae (algal turf), influenced coral recruitment, controlled invertebrates, and other herbivorous fish [18–21]. The abundance of reef fish on deeper water might be influenced by the high cover of coral community in 10 m than shallower area. The

dependency of pomacentrids to coral was studied in Lizard Island, GBR, which had the highest body mass increasing per week during their territory [22]. Total abundance of fish at a depth of 5m by 0.03 ind./m³ and 10m by 0.04 ind./m³.

The family *Pomacentridae* had the most abundant reef fish in Pulau Miang. It was 192 fish at a depth of 5m and 197 at a depth of 10m. Otherwise, the families of *Epiphidae* and *Cirrhitidae* had a lower abundance (Fig. 3). More than 340 species of damselfish (*Pomacentridae*) dwell primarily in coral reef habitats, and their colonies are the most visible in the coral reef environment [23]. *Pomacentriids* have been inhibited on coral reefs since 50 million years ago [6], and it was given a good role in coral reef community structure.

Moreover, scientists have reported that almost all living cycles of *Pomacentrids* through coral reef ecosystems such as larvae, settlement, adult, and recruitment. Based on diet, the *pomacentrids* fish were divided into three categories as herbivorous, planktivorous, and omnivorous. Research on 134 *Pomacentridae* fish revealed that these fish had six different types of food: phytoplankton, benthic algae, zooplankton, vagile invertebrate sessile invertebrate, and detritus [24].

The presence and role of herbivorous fish are very important to coral development and competition to win the competition with benthic fauna because carnivorous prey on algae resulted in coral can grow quickly [25]. *Pomacentrids* also play an active role in cleaning the waters because they feed on various detritus and plankton around the reef. Increasing nutrients in the water will increase the number of phytoplankton, and of course, too abundant amounts of plankton will interfere with the diet of corals that eat only nano-sized plankton [26]. As an omnivore, these fish also feed on animals living in the body of coral known as micro borer [27]. Too many holes produced micro borer in the body of corals will make the coral fragile and caused the weak structure of the coral. The presence of this fish turns out to be able to control the amount of micro borer on the body of the coral.

Two genera have dominated among other Pomacentridae in Pulau Miang. They were *Chromis Viridis* and *Dascyllus* sp. These species are closely related to branching corals, especially the genus *Acropora* [2]. They hide inside branches of *Acropora* coral in small colonies. Moreover, these fish also tag the branching of *Acropora* using chemical cues and use the correct signals for homing [28]. *Dascyllus* is also one of the genera that have many symbiotic corals. They live in small groups between the corals of *Acropora* and *Stylophora*. It was thought that there was a symbiotic relationship of mutualism between these fish and corals. Coral as a host protects fish and traps plankton so that fish do not have to come out of the coral to find food, while corals will be able to essential nutrients produced by such fish [29].

One hundred and twenty *Caesionidae* fish were found in Pulau Miang. *Caesionidae* was endemic fish of the Indo-Pacific and Red Sea waters [16, 30]. The family *Caesionidae* usually made a group of swimming (schooling) during prey, and this incident has been found in Pulau Miang, that schooling of *Caesoniids* fish over on observers view time. The family *Caesionidae*

preyed on zooplankton column of water with schooling strategi and sometimes clustered around divers in large numbers so that they almost cover the surrounding ocean scenery [2]. These fish are also known as fusiliers, which can move vertically from the sea surface to a maximum depth of 60 m around the coral reef ecosystem [31].




Fig. 3 Distribution of reef fish at stations at different depths

Chaetodontidae or butterflyfish are easily known due to their attractive color patterns, thin shape, and graceful swimming movements. *Chaetodontidae* is most commonly seen in large areas of living coral, usually at a depth of less than 20 m. They depend on coral reefs for shelter and their nutritional needs. Almost butterflyfish stop their foraging activity in the afternoon and hide on the reef's crevices except for the *Chaetodon lunula* that regularly eats after dark [2]. Seven species were described in this research. They were *Chaetodon lunulatus*, *C. oxycephalus*, *C. kleinii*, *C. unimaculatus*, *C. octofasciatus*, *Forcipiger flavissimus*, and *Heniochus varius*. There was no significant difference in the number of *chaetodontids* at different in five and ten m depth (Table 1; Fig. 3).

The number of *Labridae* fish found at ten m have twice the amount compared to a depth of 5 m. The most abundant from this family was *Cirrhilabrus exquisitus* (29), while the other ten species spread evenly at one individual each. *Labridae* was a large family of fish, consisting of 82 genera and 600 species of fish [32]. Most species are small, accomplishing a body length of less than 20 cm, although few achieve a mass of over 100 kg. *Labriids* occupy tropical marine and calm

waters worldwide and are most common in coral reefs, dead coral algae, sand and rubble, and algae ecosystems. *Labriids* fish have diverse feeding habits such as gastropods, small bivalves, algae, and other invertebrates [30]. This fish feeding with a group surrounds living coral.

3.2. Reef Fish Distribution by Station

Reef fish have their territory. They will occupy habitats that are considered suitable for each of them. For instance, one species needs more nutrient availability in their habitat, others prefer to choose temperate environmental waters for the recruitment and post-recruitment process, and others may need combine of those factors.

The availability of prey and interspecies also greatly influence a species's habitat presence [33]. The total abundance of fish at station 1, 2, 3 and 4 were 0.45; 0.10; 0.06 and 0.09 ind./m³ respectively. *Pomacentridae* and *Caesionidae* were the most abundant reef fish in Pulau Miang. As stated in Table 1, station one was occupied by 453 fish, station 2 was 100 fish, station 3 was 61 fish, and station 4 was 91 fish. The effect of live coral percent cover on the abundance of reef fish in

water has become an attraction of ecologists. Research on the abundance of reef fish has been conducted in Mataiva Atoll, which mentions species richness from *Chaetodontidae*, *Pomacentridae*, *Labridae*, *Scaridae*, *Acanthuridae*, and *Gobiidae* were significantly increasing once the coral cover raised to 5% [34]. The coral cover also has a pronounced influence on reef fish diversity, species composition, and abundance [35], and coral cover is known as a driver for reef fish biomass [36]. In this research, the live coral cover was 45% and stated as the highest percentage coral cover than other stations.

Table 1 Distribution of reef fish in Miang Island

Group of Fish	Family	Number of Fish			
		St. 1	St. 2	St. 3	St. 4
Indicator	<i>Chaetodontidae</i>	9	15	0	16
Major	<i>Balistidae</i>	6	0	0	1
	<i>Caesionidae</i>	120	0	0	0
	<i>Cirrhitidae</i>	1	0	0	0
	<i>Ephippidae</i>	1	0	0	0
	<i>Gobiidae</i>	0	0	1	2
	<i>Holocentridae</i>	3	0	0	0
	<i>Labridae</i>	10	8	7	5
	<i>Pomacentridae</i>	231	58	52	48
	<i>Pomachantidae</i>	1	2	0	0
	<i>Synodontidae</i>	1	0	0	1
	<i>Tetraodontidae</i>	3	0	0	0
	<i>Zanclidae</i>	10	2	0	0
	Target	<i>Acanthuridae</i>	18	4	0
<i>Lutjanidae</i>		2	7	0	0
<i>Mullidae</i>		7	0	1	4
<i>Nemipteridae</i>		7	0	0	3
<i>Serranidae</i>		11	0	0	0
<i>Siganidae</i>		1	0	0	2
Eco. Importance	<i>Scaridae</i>	11	3	0	6
Total		453	100	61	91

The number and composition of reef fish in Pulau Miang were not distributed evenly. For instance, Family *Caesionidae* was abundant in station 1 (Table 1) but not followed by three other stations. In contrast, *Pomacentridae* can be found in all stations so that the total number of this family was very high (389) (Table 1). Understanding how the pattern of reef fish distribution to their habitat is not easy because the composition of reef fish that have separated 10's meter each other has a very real difference compared to coral reef habitats that are 1000 km apart [37]. This research shows that *Caesiniidae* is only found in station one, not elsewhere, even though the distance of each station is not surely far. In another possibility, during that time number of phytoplankton in station one was very abundant and attracted *Caesionidae* to feed it. Chlorophyll contained in phytoplankton was the main food of the caesiniids fish [38]. The strong current at station one at that time and carrying much phytoplankton may have also affected the presence of these fish. The research that has been carried out is temporary. It does not study the eating habits of all types of fish for a certain period, so the habitat preferences of reef fish can only be done by making

estimates following the existing literature. Based on research conducted on Orpheus Island, Australia which states that among the many factors that affect the abundance of reef fish in a water area, the condition of corals as the main habitat for reef fish and the presence of food sources are the two most important factors that affect the diversity of reef fish [37].

4. Conclusion

Reef fish assemblage in Pulau Miang was composed of large species that *Pomacentridae* and *Caesionidae* dominated. *Pomacentridae* was very distributed on this island. They can be found easily in all stations. The habitat of these fish is mostly found in coral reef structures and eats epilithic algae in surround of coral that keep coral success on the competition with algae. *Caesionidae* is one important reef fish for human, most of the islander in Indo-Pacific has depended on these fish for protein sources. Reef fish abundance had a strong relationship with the coral reef ecosystem because the reef is the main habitat of these fish. Knowing the fish assemblages in this island accurately is important as the baseline and foundation of marine protected areas in the future. Decision-making based on ecology instead of the economy is a good approach to conserving the marine ecosystem. This study has a big effort and has involved many resources on it and if any collaboration in the future between local government, university, and private sector, the research will be expanded on revealed linkages of the environment to reef fish assemblages in this island.

Acknowledgment

We are thankful to the members of the Acropora Study Club: Achmad Rifqi Rachmatullah, Elda Pebrizayanti, and Fairly Yulianto, for their nice work on collecting data photographing both reef fish and coral in Miang Island. We are also grateful to the staff of STIPER for their assistance during the fieldwork. This work was funded in part by the Government of Kutai Timur, and all diving equipment also boat in this research was supplied by the Department of Fisheries and Marine of Kutai Timur Regency.

References

- [1] GABORIAU, T., LEPRIEUR, F., MOUILLOT, D., and HUBERT, N. Influence of the geography of speciation on current patterns of coral reef fish biodiversity across the Indo-Pacific. *Ecography*, 2018, 41(8): 1295-1306.
- [2] ALLEN, G.R., and ERDMANN, M.V. Reef Fishes of the East Indies: volumes I-III, Tropical Reef Research. *Perth, Australia*, 2012, 1(3): 1292.
- [3] TORNABENE, L., VALDEZ, S., ERDMANN, M., and PEZOLD, F. Support for a 'Center of Origin' in the Coral Triangle: Cryptic diversity, recent speciation, and local endemism in a diverse lineage of reef fishes (*Gobiidae: Eviota*). *Molecular Phylogenetics and Evolution*, 2015, 82: 200-210.
- [4] BRIGGS, J.C. Centrifugal speciation, and centers of

- origin. *Journal of Biogeography*, 2000, 27(5): 1183-1188.
- [5] WOODLAND, D.J. Zoogeography of the Siganidae (Pisces): an interpretation of distribution and richness patterns. *Bulletin of marine science*, 1983, 33(3): 713-717.
- [6] BELLWOOD, D.R., and WAINWRIGHT, P.C. The history and biogeography of fishes on coral reefs. *Coral reef fishes: dynamics and diversity in a complex ecosystem*, 2002, 532.
- [7] PELLISSIER, L. Quaternary coral reef refugia preserved fish diversity. *Science*, 2014, 344(6187): 1016-1019.
- [8] DÍAZ-PÉREZ, L. Coral reef health indices versus the biological, ecological and functional diversity of fish and coral assemblages in the Caribbean Sea. *PloS one*, 2016, 11(8): e0161812.
- [9] WILLIAMSON, D.H., CECCARELLI, D.M., EVANS, R.D., JONES, G.P., RUSS, G.R. Habitat dynamics, marine reserve status, and the decline and recovery of coral reef fish communities. *Ecology and Evolution*, 2014, 4(4): 337-354.
- [10] CECCARELLI, D.M., JONES, G.P., and MCCOOK, L.J. Interactions between herbivorous fish guilds and their influence on algal succession on a coastal coral reef. *Journal of Experimental Marine Biology and Ecology*, 2011, 399(1): 60-67.
- [11] RICE, M.M., EZZAT, L., and BURKEPILE, D.E. Corallivory in the Anthropocene: interactive effects of anthropogenic stressors and coral ivory on coral reefs. *Frontiers in Marine Science*, 2019, 5525.
- [12] VAN QUAN, N., and VINH, C.T. Study on the potential usage of butterflyfishes (*Chaetodontidae*) as bioindicators for coral reef health. *Science and Technology Journal of Agriculture and Rural Development*, 2009.
- [13] MOH'D, M.N. *The use of butterflyfishes (Chaetodontidae) as bioindicators of coral reef degradation along the coast of Tanzania*. University of Dar es Salaam, 2003. Ph.D. diss.
- [14] IRAWANSYAH I., ROSDIANTO, R., and OKTIYAS, M.L. *Terumbu Karang di Kutai Timur: Pulau Miang*. UMM Press, 2019.
- [15] ENGLISH, S.A., BAKER, V.J., and WILKINSON, C.R. *Survey manual for tropical marine resources*. Australian Institute of Marine Science Townsville, 1997.
- [16] ALLEN, G.R., and ADRIM, M. Coral reef fishes of Indonesia. *Zoological Studies-Taipei*, 2003, 42(1): 1-72.
- [17] LUTHFI, O.M., ROSDIANTO, PEBRIZAYANTI, E., ISDIANTO, A., ASADI, M.A., AFFANDI, M., and PUTRANTO, T.W.C. Prevalence and incidence of white syndrome in *Echinopora lamellosa* coral at nature reserve Pulau Sempu, Malang, Indonesia. *Ecology, Environment, and Conservation*, 2020, 26(June Suppl. Issue): S179-S185.
- [18] CECCARELLI, D.M. Modification of benthic communities by territorial damselfish: a multi-species comparison. *Coral reefs*, 2007, 26(4): 853-866.
- [19] FERREIRA, C.E.L., GONÇALVES, J.E.A., COUTINHO, R., and PERET, A.C. Herbivory by the dusky damselfish *Stegastes fuscus* (Cuvier, 1830) in a tropical rocky shore: effects on the benthic community. *Journal of Experimental Marine Biology and Ecology*, 1998, 229(2): 241-264.
- [20] WATSON, M., RIGHTON, D., AUSTIN, T., and ORMOND, R. The effects of fishing on coral reef fish abundance and diversity. *Journal of the Marine Biological Association of the United Kingdom*, 1996, 76(1): 229-233.
- [21] MENDES, T.C., QUIMBAYO, J.P., BOUTH, H.F., SILVA, L.P.S., and FERREIRA, C.E.L. The omnivorous triggerfish *Melichthys niger* is a functional herbivore on an isolated Atlantic oceanic island. *Journal of fish biology*, 2019.
- [22] DEPCZYNSKI, M., FULTON, C.J., MARNANE, M.J., and BELLWOOD, D.R. Life history patterns shape energy allocation among fishes on coral reefs. *Oecologia*, 2007, 153(1): 111-120.
- [23] FABRI, G., PARMENTIER, E., LEPOINT, G., VANDEWALLE, P., and FREDERICH, B. Diet of Damselfishes (*Pomacentridae*): a multidisciplinary approach, 2006.
- [24] FRÉDÉRICH, B., FABRI, G., LEPOINT, G., VANDEWALLE, P., and PARMENTIER, E. Trophic niches of thirteen damselfishes (*Pomacentridae*) at the Grand Récif of Toliara, Madagascar. *Ichthyological Research*, 2009, 56(1): 10-17.
- [25] CHASE, T.J., PRATCHETT, M.S., and HOOGENBOOM, M.O. Behavioral trade-offs and habitat associations of coral-dwelling damselfishes (family *Pomacentridae*). *Marine Ecology Progress Series*, 2020, 633141-156.
- [26] GRUTTER, A.S., BEJARANO, S., CHENEY, K.L., GOLDIZEN, A.W., SINCLAIR-TAYLOR, T., and WALDIE, P.A. Effects of the cleaner fish *Labroides dimidiatus* on grazing fishes and coral reef benthos. *Marine Ecology Progress Series*, 2020, 64399-114.
- [27] WEINSTEIN, D.K., MAHER, R.L., and CORREA, A.M.S. Bioerosion, in Mesophotic Coral Ecosystems. *Springer*, 2019, 829-847.
- [28] BEN-TZVI, O., TCHERNOV, D., and KIFLAWI, M. Role of coral-derived chemical cues in microhabitat selection by settling *Chromis Viridis*. *Marine Ecology Progress Series*, 2010, 409181-187.
- [29] ZIKOVA, A.V., BRITAEV, T.A., IVANENKO, V.N., and MIKHEEV, V.N. Planktonic and symbiotic organisms in the nutrition of coralobiont fish. *Journal of Ichthyology*, 2011, 51(9): 769-775.
- [30] DIBATTISTA, J.D. On the origin of endemic species in the Red Sea. *Journal of Biogeography*, 2016, 43(1): 13-30.
- [31] CARPENTER, K.E., and NIEM, V.H. *The living marine resources of the Western Central Pacific. v. 1: Seaweeds, corals, bivalves, and gastropods.-v. 2: Cephalopods, crustaceans, holothurians, and sharks.-v. 3: Batoid fishes, chimaeras, bony fishes, pt. 1 (Elopidae to Linophryniidae).-v. 4*, 1998.
- [32] WESTNEAT, M.W., and ALFARO, M.E. Phylogenetic relationships and evolutionary history of the reef fish family Labridae. *Molecular phylogenetics and evolution*, 2005, 36(2): 370-390.
- [33] BELLWOOD, D.R., STREIT, R.P., BRANDL, S.J., and TEBBETT, S.B. The meaning of the term 'function' in ecology: A coral reef perspective. *Functional Ecology*, 2019, 33(6): 948-961.
- [34] BELL, J.D., and GALZIN, R. Influence of live coral cover on coral-reef fish communities. *Marine Ecology Progress Series*, 1984, 15(3): 265-274.
- [35] KOMYAKOVA, V., MUNDAY, P.L., and JONES, G.P. Relative importance of coral cover, habitat complexity and diversity in determining the structure of reef fish communities. *PloS one*, 2013, 8(12): e83178.
- [36] RUSS, G.R., RIZZARI, J.R., ABESAMIS, R.A., and ALCALA, A.C. Coral cover a stronger driver of reef fish trophic biomass than fishing. *Ecological Applications*, 2021, 31(1): e02224.

- [37] OAKLEY-COGAN, A., TEBBETT, S.B., and BELLWOOD, D.R. Habitat zonation on coral reefs: Structural complexity, nutritional resources, and herbivorous fish distributions. *PLoS One*, 2020, 15(6): e0233498.
- [38] YAP-DEJETO, L., SIGUA, P., OSTREA, J.L.M., and RUIZO, E.K.C. Phytoplankton and Fish Abundance in the outer San Pedro Bay, Leyte, Philippines during 2013-2014. *Philippine Journal of Natural Sciences*, 2016, 21(2): 17-30.

參考文:

- [1] GABORIAU, T.、LEPRIEUR, F.、MOUILLOT, D. 和 HUBERT, N. 物種形成地理對整個印度-太平洋珊瑚礁魚類生物多樣性當前模式的影響。生態學, 2018, 41(8): 1295-1306。
- [2] 艾倫, G.R. 和 ERDMANN, M.V. 東印度群島的珊瑚礁魚類: I-III 卷, 熱帶珊瑚礁研究。澳大利亞珀斯, 2012 年, 1(3): 1292。
- [3] TORNABENE, L.、VALDEZ, S.、ERDMANN, M. 和 PEZOLD, F. 支持珊瑚三角區的“起源中心”: 隱秘多樣性、最近物種形成和不同譜系中的地方特有性礁魚(戈壁科: 埃維奧塔)。分子系統發育與進化, 2015, 82: 200-210。
- [4] BRIGGS, J.C. 離心物種形成和起源中心。生物地理學雜誌, 2000, 27(5): 1183-1188。
- [5] WOODLAND, D.J. 蠶科(雙魚座)的動物地理學: 分佈和豐富度模式的解釋。海洋科學公報, 1983, 33(3): 713-717。
- [6] BELLWOOD, D.R. 和 WAINWRIGHT, P.C. 珊瑚礁魚類的歷史和生物地理學。珊瑚礁魚類: 複雜生態系統中的動態和多樣性, 2002 年, 532。
- [7] PELLISSIER, L. 第四紀珊瑚礁保護區保護了魚類多樣性。科學, 2014, 344(6187): 1016-1019。
- [8] DÍAZ-PÉREZ, L. 珊瑚礁健康指數與加勒比海魚類和珊瑚組合的生物、生態和功能多樣性。公共廣播電台一, 2016, 11(8): e0161812。
- [9] WILLIAMSON, D.H.、CECCARELLI, D.M.、EVANS, R.D.、JONES, G.P. 和 RUSS, G.R. 棲息地動態、海洋保護區狀況以及珊瑚礁魚類群落的衰退和恢復。生態與進化, 2014, 4(4): 337-354。
- [10] CECCARELLI, D.M., JONES, G.P. 和 MCCOOK, L.J. 食草魚類行會之間的相互作用及其對沿海珊瑚礁藻類演替的影響。實驗海洋生物學與生態學雜誌, 2011, 399(1): 60-67。
- [11] RICE, M.M.、EZZAT, L. 和 BURKEPILE, D.E. 人類世的珊瑚食: 人為壓力源和珊瑚食對珊瑚礁的交互作用。海洋科學前沿, 2019, 5525。
- [12] VAN QUAN, N. 和 VINH, C.T. 研究蝴蝶魚(毛齒獸科)作為珊瑚礁健康生物指標的潛在用途。農業與農村發展科技學報, 2009。
- [13] MOHD, M.N. 使用蝴蝶魚(毛齒獸科)作為坦桑尼亞海岸珊瑚礁退化的生物指標。達累斯薩拉姆大學, 2003 年。博士論文。
- [14] IRAWANSYAH I.、ROSDIANTO, R. 和 OKTIYAS, M.L. 東庫台的珊瑚礁: 綿島。烏姆出版社, 2019 年。
- [15] ENGLISH, S.A.、BAKER, V.J. 和 WILKINSON, C.R. 熱帶海洋資源調查手冊。澳大利亞海洋科學研究所湯斯維爾, 1997。
- [16] ALLEN, G.R. 和 ADRIM, M. 印度尼西亞的珊瑚礁魚類。動物學-台北, 2003, 42(1): 1-72。
- [17] LUTHFI, O.M., ROSDIANTO, PEBRIZAYANTI, E., ISDIANTO, A., ASADI, M.A., AFFANDI, M. 和 PUTRANTO, T.W.C. 印度尼西亞瑪瑯森島自然保護區棘球蚧珊瑚中白色綜合徵的患病率和發病率。生態、環境與保育, 2020, 26(六月增刊 問題): S179-S185。
- [18] CECCARELLI, D.M. 領土雀鯛對底棲群落的改造: 多物種比較。珊瑚礁, 2007, 26(4): 853-866。
- [19] FERREIRA, C.E.L.、GONÇALVES, J.E.A.、COUTINHO, R. 和 PERET, A.C. 草食動物, 由位於熱帶岩石海岸的昏暗雀鯛 墨魚(居維葉, 1830 年) 撰寫: 對底棲群落的影響。實驗海洋生物學與生態學雜誌, 1998, 229(2): 241-264。
- [20] WATSON, M.、RIGHTON, D.、AUSTIN, T. 和 ORMOND, R. 捕魚對珊瑚礁魚類豐富度和多樣性的影響。英國海洋生物協會雜誌, 1996 年, 76(1): 229-233。
- [21] MENDES, T.C., QUIMBAYO, J.P., BOUTH, H.F., SILVA, L.P.S. 和 FERREIRA, C.E.L. 雜食性引金魚黑魚是一種生活在一個孤立的大西洋島嶼上的功能性食草動物。魚類生物學雜誌, 2019。
- [22] DEPCZYNSKI, M., FULTON, C.J., MARNANE, M.J. 和 BELLWOOD, D.R. 生活史模式塑造了珊瑚礁魚類之間的能量分配。生態學, 2007, 153(1): 111-120。
- [23] FABRI, G., PARMENTIER, E., LEPOINT, G., VANDEWALLE, P. 和 FREDERICH, B. 雀鯛(果蠅科)的飲食: 多學科方法, 2006 年。
- [24] FRÉDÉRICH, B.、FABRI, G.、LEPOINT, G.、VANDEWALLE, P. 和 PARMENTIER, E. 馬達加斯加圖利亞拉大雷西夫的十三隻雀鯛(果蠅科)的營養壁龕。魚類學研究, 2009, 56(1): 10-17。
- [25] CHASE, T.J., PRATCHETT, M.S. 和 HOOGENBOOM, M.O. 棲息於珊瑚的雀鯛(果蠅科)的行為權衡和棲息地關聯。海洋生態學進展系列, 2020, 633141-156。
- [26] GRUTTER, A.S.、BEJARANO, S.、CHENEY, K.L.、GOLDIZEN, A.W.、SINCLAIR-TAYLOR, T. 和 WALDIE, P.A. 清潔魚 唇線蟲對放牧魚類和珊瑚礁底棲動物的影響。海洋生態學進展系列, 2020, 64399-114。
- [27] WEINSTEIN, D.K., MAHER, R.L. 和 CORREA, A.M.S. 中光珊瑚生態系統中的生物侵蝕。斯普林格, 2019 年, 829-847。
- [28] BEN-TZVI, O., TCHERNOV, D., 和 KIFLAWI, M. 珊瑚衍生化學線索在微生境選擇中的作用, 通過解決 綠色銻。海洋生態學進展系列, 2010, 409181-187。
- [29] ZIKOVA, A.V., BRITAEV, T.A., IVANENKO, V.N. 和 MIKHEEV, V.N. 珊瑚魚營養中的浮游生物和共生生物。魚類學雜誌, 2011, 51(9): 769-775。
- [30] DIBATTISTA, J.D. 關於紅海特有物種的起源。生物地理學雜誌, 2016, 43(1): 13-30。
- [31] CARPENTER, K.E. 和 NIEM, V.H. 中西部太平洋的海洋生物資源。v. 1: 海藻、珊瑚、雙殼類和腹足類。-v. 2: 頭足類、甲殼類、海參和鯊魚。-v. 3: 蝙蝠魚、嵌合體和硬骨魚, pt. 1 (豹科到 亞麻科)。-v. 4, 1998。
- [32] WESTNEAT, M.W. 和 ALFARO, M.E. 珊瑚礁魚類家族的系統發育關係和進化史。分子系統發育與進化,

2005, 36(2): 370-390。

[33] BELLWOOD, D.R., STREIT, R.P., BRANDL, S.J. 和 TEBBETT, S.B. 生態學中“功能”一詞的含義：珊瑚礁視角。功能生態學, 2019, 33(6): 948-961。

[34] BELL, J.D. 和 GALZIN, R. 活珊瑚覆蓋對珊瑚礁魚類群落的影響。海洋生態學進展叢書, 1984, 15 (3) : 265-274。

[35] KOMYAKOVA, V., MUNDAY, P.L. 和 JONES, G.P. 珊瑚覆蓋、棲息地複雜性和多樣性在決定珊瑚礁魚類群落結構方面的相對重要性。公共广播电台一, 2013, 8(12) : e83178。

[36] RUSS, G.R., RIZZARI, J.R., ABESAMIS, R.A. 和 ALCALA, A.C. 珊瑚比捕魚更能驅動珊瑚魚的營養生物量。生態應用, 2021, 31(1): e02224。

[37] OAKLEY-COGAN, A. 、 TEBBETT, S.B. 和 BELLWOOD, D.R. 珊瑚礁棲息地分區：結構複雜性、營養資源和食草魚類分佈。公共广播电台一, 2020, 15(6) : e0233498。

[38] YAP-DEJETO, L., SIGUA, P., OSTREA, J.L.M. 和 RUIZO, E.K.C. 2013-2014 年菲律賓萊特市外聖佩德羅灣的浮游植物和魚類豐度。菲律賓自然科學雜誌, 2016, 21(2): 17-30。